

Exercise capacity as a predictor of survival among ambulatory patients with end-stage renal disease

KATHY E. SIETSEMA, ANTONINO AMATO, SHARON G. ADLER, and ERIC P. BRASS

Harbor-UCLA Research and Education Institute; and Sigma Tau Research, Inc.

Exercise capacity as a predictor of survival among ambulatory patients with end-stage renal disease.

Background. Exercise capacity is reduced in end-stage renal disease (ESRD). Exercise requires the integrated function of multiple vital organs, and low exercise capacity is an independent predictor of mortality in a number of clinical populations. We analyzed the value of exercise capacity, characterized as peak oxygen uptake (VO_2), for predicting survival in a cohort of 175 hemodialysis patients over a median follow-up of 39 months.

Methods. Survival status was determined for 175 ESRD patients who had participated in previous studies for which peak VO_2 and other clinical data had been determined. Chi-square and Kaplan-Meier survival analyses were performed, and a minimal model of factors related to mortality was developed by Cox multiple regression.

Results. There were 23 deaths during the follow-up period. Peak VO_2 (>17.5 mL/min/kg) was a powerful predictor of survival ($P = 0.009$ by Kaplan-Meier). Age (<65 years), dialysis vintage (<39 months), pulse pressure (<54 mm Hg), and absence of diagnoses of diabetes or heart failure were also associated with better survival on univariate analyses. On multivariate analysis peak VO_2 contributed significantly to the minimal explanatory model relating clinical variables to mortality (overall $\chi^2 = 25.5$, $P = 0.00001$).

Conclusion. Among these ambulatory ESRD patients, peak VO_2 was a stronger predictor of survival than many traditional prognostic variables, some of which are subject to ceiling effects. Exercise capacity may thus provide incremental prognostic information concerning healthier ESRD patients. Because peak VO_2 may be modified by exercise training, the potential of exercise as an intervention to improve survival is suggested.

Many variables have been identified as predictive of mortality among patients with end-stage renal disease (ESRD). These include laboratory [1–5] and anthropometric [6] variables reflecting nutritional state, the presence of chronic inflammation [7, 8], the burden of

comorbid or complicating medical conditions [9, 10], and cardiovascular function [11–14]. Exercise capacity, characterized by maximal or peak exercise oxygen uptake (VO_2), is influenced by many of the same factors as are reflected in the previously characterized predictive variables in ESRD. As a group, patients with ESRD have marked reductions in peak VO_2 , reflecting limited capacity for physical work [15, 16]. In addition to characterizing functional capability, however, peak VO_2 has been identified as an independent predictor of mortality or survival in a number of chronic diseases [17–19]. Thus, in this analysis we sought to test the hypothesis that peak VO_2 would predict survival among patients receiving chronic hemodialysis for ESRD. Outcome data were collected for hemodialysis patients whose baseline exercise capacity and clinical status were carefully assessed during their participation in two completed clinical trials, and the relationship of peak VO_2 to survival was evaluated.

METHODS

Study subjects

The analysis made use of a database of clinical, demographic, and exercise variables for patients with ESRD who participated in two clinical trials of L-Carnitine (Carnitor[®]) supplementation in chronic hemodialysis (CHIEF studies) [20]. Key inclusion criteria for those studies included ESRD on thrice weekly hemodialysis for at least 6 months with Kt/V value of ≥ 1.2 , clinical stability, and ability to perform incremental exercise testing to symptomatic maximum. Study participation was six months' duration, during which time peak VO_2 was determined at three-month intervals using a standardized cycle ergometry exercise testing protocol. Patients were excluded from study if they could not cooperate with testing or if an initial screening exercise test had to be terminated because of adverse reactions such as ischemic changes or arrhythmia on electrocardiogram, or pathologic blood pressure response. The present analysis is based on peak VO_2 measures determined on a baseline test performed on a separate day from the screening test.

Key words: oxygen uptake, VO_2 , exercise, mortality, end-stage renal disease, hemodialysis.

Received for publication June 30, 2003
and in revised for September 8, 2003
Accepted for publication September 22, 2003

© 2004 by the International Society of Nephrology

Table 1. Patient characteristics

Variable	Units	All patients	Survivors	Non-survivors	P value ^a
N		175	152	23	
Age	years	46 ± 13	44 ± 12	56 ± 16	0.000074
Gender	% male	65	65	61	0.69
Peak VO ₂	mL/min/kg	18.6 ± 6.5	19.2 ± 6.6	14.9 ± 4.9	0.003
Dialysis vintage	months	41.5 ± 15	40 ± 13	52 ± 18	0.0002
Serum albumin	g/dL	3.97 ± 0.28	3.99 ± 0.27	3.87 ± 0.30	0.06
Serum creatinine	mg/dL	10.9 ± 2.9	11.2 ± 2.8	9.6 ± 3.2	0.02
Blood pressure, resting					
Systolic	mm Hg	138 ± 23	136 ± 23	149 ± 22	0.01
Diastolic	mm Hg	82 ± 13	83 ± 13	80 ± 13	0.37
Pulse pressure	mm Hg	55 ± 19	53 ± 18	68 ± 20	0.0003
Hemoglobin	g/dL	11.2 ± 1.1	11.3 ± 1.1	10.9 ± 1.0	0.11
Race	% non-white	69	71	52	0.60
Diabetes	%	19	15	44	0.001
CHF	%	10	7	26	0.004
BMI	kg/m ²	24.5 ± 4.1	24.5 ± 4.1	24.7 ± 4.3	0.87

Abbreviations are: CHF, chronic heart failure; pulse pressure, systolic pressure minus diastolic pressure; BMI, body mass index. Values are mean ± 1 SD.

^aFor differences between survivors and non-survivors [by analysis of variance (ANOVA) for continuous variables and χ^2 for categorical variables].

Approximately three years after completion of the CHIEF studies, a questionnaire was mailed to each participating site to ascertain survival status and dates and causes of deaths as of October 1, 2001 for patients who had been enrolled in the trials. Ten of the original 12 study sites participated in the follow-up. Collection and use of these data for the present analysis was reviewed and approved by the institutional review board of the Harbor-UCLA Research and Education Institute.

Analyses

Patient characteristics as related to outcomes were summarized using descriptive statistics. Group values are reported as mean and SD unless otherwise noted. The primary outcome measure was survival during the follow-up period. Simple descriptive statistics were used to characterize survivors and non-survivors, who were compared using either analysis of variance (ANOVA) or chi-square analysis with Yate's correction where applicable. Survival status was analyzed as a function of exercise capacity, expressed as peak VO₂ in mL/kg/min, and of other selected clinical variables. Chi-square analysis was used to compare crude death rates and univariate survival analyses were performed using the log-rank test on survival curves created with the method of Kaplan-Meier. For these purposes stratification of the population for continuous variables was done using values previously utilized in survival analyses for this patient population (for age and serum albumin), or using the observed group median value (for peak VO₂, dialysis vintage, serum creatinine, blood pressures, and hemoglobin). Sensitivity analyses were also conducted for some of these variables, using different cut-off points. Survival data were not censored at the time of kidney transplantation. To control for covariance of exercise capacity with other variables recognized to have prognostic significance in this population,

Cox's proportional-hazards multivariate regression was performed. All potentially relevant variables identified from Kaplan-Meier survival analyses ($P < 0.15$) were entered into the Cox model; a minimal explanatory model predicting mortality was then derived using a backward selection approach. All analyses were performed using Statistica v. 5.1 software (StatSoft, Inc., Tulsa, OK, USA).

RESULTS

Data were available for all patients from the 10 reporting study sites, representing 175 of 193 patients enrolled in the CHIEF studies. Survival status and dates of deaths were determined approximately 3.5 years after the initial baseline assessments were performed (mean follow up: 1242 ± 273 days). There were a total of 23 deaths (13%) during the follow-up period. Cardiovascular causes were reported for seven deaths (including cardiac arrest, cardiomyopathy, and cerebral ischemia), infectious causes for seven (including sepsis, abscess, and pneumonia), and malignancy for two. Causes of the remaining seven deaths were reported as unknown. Forty-four patients received kidney transplants during the follow-up period, and two of the deaths occurred among these patients.

Subject characteristics

Baseline characteristics for all patients and for the subgroups of surviving and non-surviving patients are shown in Table 1. Patients averaged 46 ± 13 years of age and had been on dialysis an average of 41 ± 15 months at the time of the baseline evaluations. Group mean values for clinical variables reflect, in part, the entry criteria for the original study which included requirements of clinical stability, adequate dialysis as reflected in a Kt/V of at least 1.2, and treatment with erythropoietin in accordance with prevailing standards of clinical practice.

Table 2. Crude mortality rates for 175 patients with ESRD stratified according to selected clinical variables

Variable	Stratification	Deaths/group (%)	total	<i>P</i> value ^a
Age years	<65	17/157 (11)	175	0.02
	≥65	6/18 (33)		
Peak VO ₂ mL/min/kg	≤17.5	19/88 (22)	175	0.002
	>17.5	4/87 (5)		
Dialysis vintage months	≤39	6/88 (7)	175	0.02
	>39	17/87 (20)		
Serum albumin g/dL	<4.0	14/75 (19)	173 ^b	0.15
	≥4.0	9/98 (9)		
Serum creatinine mg/dL	≤10.8	16/90 (18)	175	0.17
	>10.8	7/85 (8)		
Systolic blood pressure mm Hg	≤138	7/88 (8)	175	0.07
	>138	16/87 (18)		
Pulse pressure mm Hg	≤54	6/88 (7)	175	0.02
	>54	17/87 (20)		
Hemoglobin g/dL	≤11.3	14/86 (16)	171 ^b	0.38
	>11.3	9/85 (11)		
Diagnosis of CHF	Yes	6/17 (35)	175	0.01
	No	17/158 (11)		
Diagnosis of diabetes	Yes	10/33 (30)	175	0.003
	No	13/142 (9)		

Abbreviations are: pulse pressure, systolic pressure minus diastolic pressure; CHF, chronic heart failure.

^aBy Yates' χ^2 test.

^bDue to missing values (albumin, 2; hemoglobin, 4).




Fig. 1. Survival as a function of baseline value of peak exercise VO₂ for 175 ambulatory patients with end-stage renal disease. Patients with peak VO₂ above the median value of 17.5 mL/min/kg (interrupted line) had significantly better survival than those with lower values (solid line) (*P* = 0.009 by log-rank test).




Fig. 2. Survival as a function of baseline serum albumin for 175 ambulatory patients with end-stage renal disease. Serum albumin concentration at or above the group median value of 4.0 gm/dL (interrupted line) showed a trend toward better survival than those with lower values (solid line) (*P* = 0.08 by log-rank test).

Factors influencing survival

Crude death rates for patients stratified by selected clinical variables are shown in Table 2. Patients with a peak VO₂ above the group median value of 17.5 mL/min/kg had a significantly better crude survival rate than those with lower peak VO₂ (Table 2); this finding was confirmed by univariate Kaplan-Meier analysis (*P* = 0.009) (Fig. 1). Similar results were obtained if peak VO₂ was stratified at 14 or 15 mL/min/kg (*P* = 0.0016 and 0.00064, respectively; data not shown). Patients younger than age 65, those with resting pulse pressure of less than 54, and those without a diagnosis of diabetes, or without a diagnosis of chronic heart failure, had a better crude

survival than those who were older, had a higher pulse pressure, were diabetic, or suffered from chronic heart failure. There was a trend (*P* = 0.15) toward better crude survival for those whose serum albumin was at or above the group median value of 4.0 g/dL (Table 2), and this trend was confirmed on univariate Kaplan-Meier analysis (*P* = 0.08) (Fig. 2). In contrast, hemoglobin and serum creatinine were not significantly related to mortality in this analysis (Table 2), nor were race or gender (data not shown).

Multivariate analysis was somewhat limited by the relatively low number of deaths and the large number of variables available. Nonetheless, a highly significant

Table 3. Independent variables and minimal model of factors related to mortality in 175 patients with ESRD

Variable	Initial variables		Minimal model	
	Beta	P value	Beta	P value
Age years	0.04	0.40	0.05	0.002
Peak VO ₂ mL/min/kg	-0.09	0.17	-0.001	0.05
Diabetes yes	0.06	0.90	—	—
CHF yes	1.12	0.04	1.04	0.03
Hb g/dL	-0.06	0.80	—	—
Albumin g/dL	-0.24	0.78	—	—
Systolic blood pressure mm Hg	0.00	0.99	—	—
Pulse pressure mm Hg	0.02	0.32	—	—
Dialysis vintage months	-0.01	0.84	—	—
Creatinine mg/dL	-0.08	0.40	—	—

Abbreviations are: pulse pressure, systolic pressure minus diastolic pressure; CHF, chronic heart failure. Initial variables were those found to be related to survival on Kaplan-Meier analysis at $P < 0.15$.

minimal model (overall $\chi^2 = 25.5$, $P = 0.00001$) was developed that included age, peak VO₂, and the diagnosis of chronic heart failure as predictors of mortality (Table 3).

DISCUSSION

In this analysis of data from 175 ambulatory hemodialysis patients, peak VO₂ was a powerful predictor of survival status over an average follow-up of approximately 3.5 years. In a multivariate minimal explanatory model, the only additional factors significantly enhancing this predictive value were age and the presence or absence of chronic heart failure.

Individual variables related to prognosis in hemodialysis populations include functions of nutrition, inflammation, cardiovascular status, and comorbid medical conditions. Serum concentrations of albumin [1], prealbumin [2], and creatinine [1] all have positive relationships to survival, consistent with their relationship to protein balance and nutritional state. There are negative associations between inflammatory markers and survival [8, 21], which may relate in part to the negative effect of inflammation on protein balance [22] as well as pathogenic processes underlying vascular disease [7]. More distal measures of cardiovascular dysfunction, including left ventricular hypertrophy [14] and pulse pressure [12], or chronic heart failure, are associated with mortality, as is the diagnoses of diabetes [10, 23]. Not surprisingly, many of these factors covary, reflecting both commonality of cause and physiologic interactions among the processes. Consistent with this, in the present analysis a number of variables including serum albumin, serum creatinine, and dialysis vintage, which were each significantly related to outcome on univariate analysis, were effectively incorporated into a smaller number of variables on multivariate regression.

Exercise capacity of healthy persons is intrinsically dependent on many of the same variables that predict survival in hemodialysis patients [24, 25]. Specifically, the

capacity to transport oxygen is determined by cardiovascular function and hemoglobin concentration, and the capacity for utilization of oxygen is related to skeletal muscle mass and function, which are dependent on nutritional status. Both diabetes and hypertension, which may predict poor outcome in ESRD, are associated with reduced exercise capacity even in persons without renal failure [26, 27]. We have previously reported correlations between exercise capacity of hemodialysis patients and a number of these variables, including hemoglobin concentration, serum albumin, and the diagnosis of diabetes [28]. It is thus reasonable to expect that peak exercise VO₂ would represent an integrated measure of multiple important prognostic variables in ESRD, and therefore would itself be related to outcome. Apart from serving indirectly as a marker of other prognostic factors, however, peak VO₂ directly reflects the capacity to increase cardiac output in response to the physiologic stress of exercise. As such, it is likely to correspond to the capacity to survive pathologic stresses related to acute illness or injury, as well.

The patients included in this analysis represent a healthy subset of the hemodialysis population, reflected in an overall mortality rate that was far lower than that reported for United States hemodialysis patients in general [29]. By initial study inclusion criteria, the patients were judged to be clinically stable and able to perform maximal exercise testing. Non-ambulatory, frail, or unstable patients, who would be predicted to be at highest risk for mortality based on commonly used prognostic variables, are, thus, not represented. This may explain why serum albumin was a relatively weak predictor of survival in this study, despite being among the strongest reported prognostic variables in large ESRD cohorts [1]. Patients who had findings of significant exercise-induced EKG abnormalities on screening exercise tests were also excluded from the study, although a substantial proportion of the observed mortality was nevertheless attributed to cardiovascular disease. The prognostic power of peak VO₂ in this relatively healthy subset of the dialysis population is thus of particular interest, as the mortality risk of these patients may be less readily predicted by other assessments due to threshold effects [30].

Exercise capacity has been recognized as an independent predictor of survival across a broad range of values within healthy populations [31–34]. A similar relationship between exercise capacity and prognosis has been identified in certain clinical conditions, particularly cardiovascular diseases [19, 35]. This is particularly well-documented in chronic heart failure [17], and has practical application in prioritization and timing of heart transplantation. In that setting, peak VO₂ values less than 14 mL/min/kg have been used as a criteria for heart transplant eligibility [36], based on observations that higher values predicted a two-year survival as good as that of

coincident new transplant recipients. In neither the heart failure population [17] nor among apparently healthy men [34] does there appear to be a ceiling above which the relationship between peak VO₂ and prognosis is lost, even after control for other commonly used clinical and demographic risk factors.

The number of deaths in the present study was low, limiting the statistical power of multivariate analyses to fully explore the relationships between peak VO₂, other clinical factors, and mortality. The highly significant relationship between exercise capacity and survival on univariate analysis, and its contribution to the multivariate model supports further investigation of this, however, particularly with respect to risk stratification of the healthiest and most functional ESRD patients.

Ultimately, the value of identifying predictors of survival or mortality lies in the identification of modifiable factors to target for intervention. It is noteworthy therefore that peak VO₂ may be increased by physical exercise training. Furthermore, it appears that training reduces mortality in the general population [37, 38] and perhaps among patients with heart failure [39]. Among patients with ESRD exercise training has been demonstrated to increase peak VO₂ [40] and functional capacity [41]. It has also recently been reported [42] that dialysis patients who identify themselves as sedentary have higher mortality than those who report being ambulatory. Whether sedentary behavior contributes causally to increased mortality in this setting, or is simply a marker of more extensive disease and poorer prognosis is not clear. Thus, it remains to be seen whether increasing peak VO₂ by exercise training might modify mortality rate in this population.

CONCLUSION

In this study peak VO₂ was found to be a highly significant predictor of survival among a stable ambulatory cohort of hemodialysis patients. This finding is consistent with other data demonstrating that nutritional status, cardiovascular factors, and exercise behavior are significant factors related to outcome in ESRD, and suggests that exercise testing may represent a useful approach to risk stratification among the most functional patients with ESRD. The potential for interventions targeting exercise function to further enhance survival in this population is deserving of further study.

ACKNOWLEDGMENTS

The authors wish to thank the investigators of the CHIEF trials for providing data for this analysis. This project was supported in part by Sigma Tau Research, Inc.

Reprint requests to Kathy E. Sietsema, M.D., Harbor-UCLA Research and Education Institute, 1124 West Carson Street, Torrance, CA 90502. E-mail: ksietsema@rei.edu

REFERENCES

1. LOWRIE EG, LEW NL: Death risk in hemodialysis patients: The predictive value of commonly measured variables and an evaluation of death rate differences between facilities. *Am J Kidney Dis* 15:458–482, 1990
2. MITTMAN N, AVRAM MM, OO KK, CHATTOPADHYAY J: Serum prealbumin predicts survival in hemodialysis and peritoneal dialysis: 10 years of prospective observation. *Am J Kidney Dis* 38:1358–1364, 2001
3. AVRAM MM, MITTMAN N, BONOMINI L, et al: Markers for survival in dialysis: a seven-year prospective study. *Am J Kidney Dis* 26:209–219, 1995
4. COMBE C, CHAUVEAU P, LAVILLE M, et al: Influence of nutritional factors and hemodialysis adequacy on the survival of 1,610 French patients. *Am J Kidney Dis* 37:S81–S88, 2001
5. LEAVEY SF, STRAWDERMAN RL, JONES CA, et al: Simple nutritional indicators as independent predictors of mortality in hemodialysis patients. *Am J Kidney Dis* 31:997–1006, 1998
6. KOPPLE JD, ZHU X, LEW NL, LOWRIE EG: Body weight-for-height relationships predict mortality in maintenance hemodialysis patients. *Kidney Int* 56:1136–1148, 1999
7. ZIMMERMANN J, HERRLINGER S, PRUY A, et al: Inflammation enhances cardiovascular risk and mortality in hemodialysis patients. *Kidney Int* 55:648–658, 1999
8. YEUN JY, LEVINE RA, MANTADILOK V, KAYSAN GA: C-Reactive protein predicts all-cause and cardiovascular mortality in hemodialysis patients. *Am J Kidney Dis* 35:469–476, 2000
9. MAILLOUX LU, NAPOLITANO B, BELLUCCI AG, et al: The impact of co-morbid risk factors at the start of dialysis upon the survival of ESRD patients. *ASAIO J* 42:164–169, 1996
10. Comorbid conditions and correlations with mortality risk among 3,399 incident hemodialysis patients. *Am J Kidney Dis* 20:32–38, 1992
11. BLACHER J, PANNIER B, GUERIN AP, et al: Carotid arterial stiffness as a predictor of cardiovascular and all-cause mortality in end-stage renal disease. *Hypertension* 32:570–574, 1998
12. KLASSEN PS, LOWRIE EG, REDDAN DN, et al: Association between pulse pressure and mortality in patients undergoing maintenance hemodialysis. *JAMA* 287:1548–1555, 2002
13. SAFAR ME, BLACHER J, PANNIER B, et al: Central pulse pressure and mortality in end-stage renal disease. *Hypertension* 39:735–738, 2002
14. SILBERBERG JS, BARRE PE, PRICHARD SS, SNIDERMAN AD: Impact of left ventricular hypertrophy on survival in end-stage renal disease. *Kidney Int* 36:286–290, 1989
15. PAINTER P, MESSER-REHAK D, HANSON P, et al: Exercise capacity in hemodialysis, CAPD, and renal transplant patients. *Nephron* 42:47–51, 1986
16. JOHANSEN KL: Physical functioning and exercise capacity in patients on dialysis. *Adv Ren Replace Ther* 6:141–148, 1999
17. MYERS J, GULLESTAD L, VAGELOS R, et al: Clinical, hemodynamic, and cardiopulmonary exercise test determinants of survival in patients referred for evaluation of heart failure. *Ann Intern Med* 129:286–293, 1998
18. NIXON PA, ORENSTEIN DM, KELSEY SF, DOERSHUK CF: The prognostic value of exercise testing in patients with cystic fibrosis. *N Engl J Med* 327:1785–1788, 1992
19. VANHEES L, FAGARD R, THIJS L, et al: Prognostic significance of peak exercise capacity in patients with coronary artery disease. *J Am Coll Cardiol* 23:358–363, 1994
20. BRASS EP, ADLER S, SIETSEMA KE, et al, for the CHIEF investigators: Effect of intravenous l-carnitine supplementation on carnitine concentrations, maximal exercise capacity and quality of life in maintenance hemodialysis patients. *Am J Kidney Dis* 37:1018–1028, 2001
21. KAYSAN GA: The microinflammatory state in uremia: causes and potential consequences. *J Am Soc Nephrol* 12:1549–1557, 2001
22. OWEN WF, LOWRIE EG: C-reactive protein as an outcome predictor for maintenance hemodialysis patients. *Kidney Int* 54:627–636, 1998
23. LOWRIE EG, LEW NL, HUANG WH: Race and diabetes as death risk predictors in hemodialysis patients. *Kidney Int (Suppl)* 38:S22–S31, 1992

24. JONES NL, KILLIAN KJ: Exercise limitation in health and disease. *N Engl J Med* 343:632–641, 2000
25. WASSERMAN K, WHIPP BJ: Exercise physiology in health and disease (state of the art). *Am Rev Respir Dis* 112:219–249, 1975
26. ESTACIO RO, REGENSTEINER JG, WOLFEL EE, et al: The association between diabetic complications and exercise capacity in NIDDM patients. *Diabetes Care* 21:291–5, 1998
27. LIM PO, MACFADYEN RJ, CLARKSON PB, MACDONALD TM: Impaired exercise tolerance in hypertensive patients. *Ann Intern Med* 124:41–55, 1996
28. SIETSEMA KE, HIATT WR, ESLER A, et al: Clinical and demographic predictors of exercise capacity in end-stage renal disease. *Am J Kidney Dis* 39:76–85, 2002
29. US Renal Data System: *USRDS 2001 Annual Data Report*. Atlas of end stage renal disease in the United States, Bethesda, MD, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, 2001
30. GOTCH FA, LEVIN NW, PORT FK, et al: Clinical outcome relative to the dose of dialysis is not what you think: The fallacy of the mean. *Am J Kidney Dis* 30:1–15, 1997
31. EKELUND LG, HASKELL WL, JOHNSON JL, et al: Physical fitness as a predictor of cardiovascular mortality in asymptomatic North American men. The Lipid Research Clinics Mortality Follow-up Study. *N Engl J Med* 319:1379–1384, 1988
32. BLAIR SN, KOHL HW, III, PAFFENBARGER RS, JR, et al: Physical fitness and all-cause mortality. A prospective study of healthy men and women. *JAMA* 262:2395–2401, 1989
33. LAUKKANEN JA, LAKKA TA, RAURAMAA R, et al: Cardiovascular fitness as a predictor of mortality in men. *Arch Intern Med* 161:825–831, 2001
34. MYERS J, PRAKASH M, FROELICHER V, et al: Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 346:793–801, 2002
35. CHURCH TS, KAMPERT JB, GIBBONS LW, et al: Usefulness of cardiorespiratory fitness as a predictor of all-cause and cardiovascular disease mortality in men with systemic hypertension. *Am J Cardiol* 88:651–656, 2001
36. MANCINI DM, EISEN H, KUSSMAUL W, et al: Value of peak exercise oxygen consumption for optimal timing of cardiac transplantation in ambulatory patients with heart failure. *Circulation* 83:778–86., 1991
37. ERIKSSON G, LIESTOL K, BJORNHOLT J, et al: Changes in physical fitness and changes in mortality. *Lancet* 352:759–762, 1998
38. BLAIR SN, KOHL HW, III, BARLOW CE, et al: Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *JAMA* 273:1093–1098, 1995
39. BELARDINELLI R, GEORGIU D, CIANCI G, PURCARO A: Randomized, controlled trial of long-term moderate exercise training in chronic heart failure: effects on functional capacity, quality of life, and clinical outcome. *Circulation* 99:1173–1182, 1999
40. PAINTER PL, NELSON-WOREL JN, HILL MM, et al: Effects of exercise training during hemodialysis. *Nephron* 43:87–92, 1986
41. PAINTER P, MOORE G, CARLSON L, et al: Effects of exercise training plus normalization of hematocrit on exercise capacity and health-related quality of life. *Am J Kidney Dis* 39:257–265, 2002
42. O'HARE AM, TAWNEY K, BACCHETTI P, JOHANSEN KL: Decreased survival among sedentary patients undergoing dialysis: Results from the dialysis morbidity and mortality study wave 2. *Am J Kidney Dis* 41:447–454, 2003